

Massive T-duality in six dimensions

Bert Janssen¹,

*Department of Mathematical Sciences
South Road, Durham DH1 3LE
United Kingdom*

ABSTRACT

A massive version of T-duality in six dimensions is given, that maps the $K3$ compactification of Romans' theory onto the $K3$ compactification of Type IIB theory. This is done by performing a (standard) Kaluza-Klein reduction on six-dimensional massive Type IIA and a Scherk-Schwarz reduction on Type IIB, mapping both theories onto the same five-dimensional theory. We also comment shortly on the difficulties arising if one intends to construct a massive generalisation of the six-dimensional string-string duality.

¹E-mail address: bert.janssen@durham.ac.uk

1 Introduction

It is generally accepted that the various string theories and their supergravity limits are all interrelated through T-, S-, or U-dualities, being different perturbative expansions in different points in the moduli space of the same M-theory (see for instance [1, 2]). Many of these duality transformations have been explicitly constructed and most of them, although not rigorously proven, are believed to hold.

There is, however, a class of supergravity theories that is much less studied than other theories, namely the massive supergravities. These are supergravities with one or more constant parameter with dimension of mass that can be interpreted as the Hodge dual of a $(D - 1)$ -form potential in D dimensions [3, 4]. Typically, massive gauge transformations give masses to some of the gauge fields by eating other fields.

Probably the best known massive supergravity theory is ten-dimensional massive Type IIA or Romans' theory [5], which is a generalisation of standard Type IIA, adding a cosmological constant term. But there are also other ways to obtain massive supergravities. Performing a generalised Scherk-Schwarz reduction [6] gives supergravity theories with mass parameters, provided that one gives extra dependences on the compactified coordinates to axionic fields. Alternatively, gauging some of the global symmetry transformations (i.e. giving them a linear dependence on the coordinates), either before or after dimensional reduction, leads to similar results [4, 7]-[15].

Some work has been done on the generalisations of the duality symmetries to massive supergravity theories. In [4] it was shown that the T-duality between ten-dimensional Type IIA and Type IIB can be extended to a massive T-duality between Romans' theory and Type IIB, linking the Kaluza-Klein reduction of Romans' theory to the Scherk-Schwarz reduction of Type IIB. It was also shown [11, 13] that the Scherk-Schwarz reduction of the Heterotic theory on T^d can be written in an $O(d, d + 16)$ symmetric way and recently an explicitly $O(4, 20)$ invariant formulation of the $K3$ compactification of Romans' theory was given [15]. The latter results therefore indicate that these massive supergravity theories have the same T-duality groups as their massless counterparts, provided that the masses also transform under these symmetry groups.

In this paper we want to make use of the results of [15] and construct the massive T-duality between the $K3$ compactification of Romans' theory and the $K3$ compactification of Type IIB supergravity, using analogous techniques as in [4]. We will gauge a particular subgroup of the $O(5, 21)$ symmetry group of Type IIB, that acts on 24 scalars and 26 vectors. Since the 24 scalars appear as axions (i.e. only via their derivative), Scherk-Schwarz reduction will give rise in five dimensions to the same number of masses and the same massive gauge transformations as one obtains through standard Kaluza-Klein reduction of six-dimensional massive Type IIA.

The paper is organised as follows: in section 2 we perform a standard Kaluza-Klein reduction of the massive Type IIA over $K3$, leading to an $O(4, 20)$ invariant massive $N = 2$ supergravity in five dimensions. In section 3 we review some of the properties of Type IIB compactified over $K3$. The Scherk-Schwarz reduction of Type IIB and the obtained massive T-duality rules are given in sections 4 and 5 and in section 6 we comment briefly on the impossibility of generalising the six-dimensional string-string duality between Type IIA and Heterotic to the massive case. In section 7 we summarise our conclusions.

2 Kaluza-Klein reduction of $m\text{IIA}/K3$

The $K3$ compactification of ten-dimensional Type IIA supergravity has been known for a long time [16, 17]. Its bosonic field content consists of a six-dimensional metric $\hat{g}_{\hat{\mu}\hat{\nu}}$, a dilaton $\hat{\phi}$, a Kalb-Ramond field $\hat{B}_{\hat{\mu}\hat{\nu}}$, 24 vectors $\hat{V}_{\hat{\mu}}^a$ forming a vector representation of $O(4, 20)$ and 80 scalars which parametrise an $O(4, 20)/(O(4) \times O(20))$ coset and can be combined into a symmetric $O(4, 20)$ matrix $\hat{M}_{ab}^{-1}$. This field content arises naturally from the geometrical properties of the $K3$ manifold, more precisely from the 22 non-contractable two-cycles and the four-cycle over which the various ten-dimensional fields can be wrapped.

Recently also a $K3$ compactification of massive ten-dimensional Type IIA supergravity has been given [15]. Here it was shown that, apart from the fields mentioned above, one also generates 24 mass parameters m^a , that transform in the vector representation of $O(4, 20)$.

Twenty two of these masses come from wrapping the ten-dimensional RR two-form field strength over the 22 two-cycles of $K3$ and one mass comes from wrapping the ten-dimensional RR four-form field strength over the four-cycle, in a kind of Scherk-Schwarz-like reduction over $K3$. This is possible because of the fact that the RR one- and three-form fields only occur via their respective field strengths (up to field redefinitions) [9]. The last mass parameter is coming directly from the mass in ten-dimensional Romans' theory. It is remarkable that one has to take into account the mass coming from Romans' theory in order to construct a full $O(4, 20)$ invariant massive Type IIA theory in six dimensions. Indeed, it is claimed in [15] that the (Scherk-Schwarz) compactification of Romans' theory over $K3$ yields a unification of various six-dimensional massive supergravity theories, related by $O(4, 20)$ symmetry.

The bosonic part of the six-dimensional massive Type IIA action ($m\text{IIA}$) is given by [15]:

$$\begin{aligned} \mathcal{L}_{m\text{IIA}} = & \sqrt{|\hat{g}|} \left\{ e^{-2\hat{\phi}} \left[\hat{R} - 4(\partial\hat{\phi})^2 + \frac{1}{12} \hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}}(\hat{B}) \hat{H}^{\hat{\mu}\hat{\nu}\hat{\rho}}(\hat{B}) + \frac{1}{8} \text{Tr}(\partial\hat{M}^{-1}\partial\hat{M}) \right] \right. \\ & \left. - \frac{1}{4} \hat{F}_{\hat{\mu}\hat{\nu}}^a(\hat{V}) \hat{M}_{ab}^{-1} \hat{F}^{b\hat{\mu}\hat{\nu}}(\hat{V}) - 2m^a \hat{M}_{ab}^{-1} m^b \right\} \\ & + \frac{1}{16} \epsilon^{\hat{\mu}\hat{\nu}\hat{\rho}\hat{\lambda}\hat{\sigma}\hat{\tau}} \eta_{ab} \left\{ \hat{F}_{\hat{\mu}\hat{\nu}}^a(\hat{V}) \hat{F}_{\hat{\rho}\hat{\lambda}}^b(\hat{V}) \hat{B}_{\hat{\sigma}\hat{\tau}} + 2m^a \hat{F}_{\hat{\mu}\hat{\nu}}^b(\hat{V}) \hat{B}_{\hat{\rho}\hat{\lambda}} \hat{B}_{\hat{\sigma}\hat{\tau}} + \frac{4}{3} m^a m^b \hat{B}_{\hat{\mu}\hat{\nu}} \hat{B}_{\hat{\rho}\hat{\lambda}} \hat{B}_{\hat{\sigma}\hat{\tau}} \right\} . \end{aligned} \quad (2.1)$$

The field strengths of the various gauge fields are defined as

$$\hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}}(\hat{B}) = 3\partial_{[\hat{\mu}} \hat{B}_{\hat{\nu}\hat{\rho}]} , \quad \hat{F}_{\hat{\mu}\hat{\nu}}^a(\hat{V}) = 2\partial_{[\hat{\mu}} \hat{V}_{\hat{\nu}]}^a + 2m^a \hat{B}_{\hat{\mu}\hat{\nu}} , \quad (2.2)$$

and η_{ab} is the invariant $O(4, 20)$ metric

$$\eta_{ab} = \begin{pmatrix} 0 & \mathbb{1}_4 & 0 \\ \mathbb{1}_4 & 0 & 0 \\ 0 & 0 & \mathbb{1}_{16} \end{pmatrix} . \quad (2.3)$$

The gauge fields transform under the following gauge transformations

$$\delta \hat{B}_{\hat{\mu}\hat{\nu}} = \partial_{[\hat{\mu}} \hat{\Sigma}_{\hat{\nu}]} , \quad \delta \hat{V}_{\hat{\mu}}^a = \partial_{\hat{\mu}} \hat{\chi}^a - m^a \hat{\Sigma}_{\hat{\mu}} . \quad (2.4)$$

Note that the $\hat{V}_{\hat{\mu}}^a$ transform as a Stückelberg field under $\hat{\Sigma}_{\hat{\mu}}$. Hence, by an appropriate redefinition of $\hat{B}_{\hat{\mu}\hat{\nu}}$,

$$m^a \hat{B}_{\hat{\mu}\hat{\nu}} = m^a \hat{B}'_{\hat{\mu}\hat{\nu}} - \partial_{[\hat{\mu}} \hat{V}_{\hat{\nu}]}^a , \quad (2.5)$$

the Kalb-Ramond field $\hat{B}'_{\hat{\mu}\hat{\nu}}$ becomes massive by eating the vectors.

The dimensional reduction to five dimensions is straight forward. Calling the direction over which we reduce x and using the standard reduction rules²

$$\begin{aligned}
\hat{g}_{xx} &= -k^2, & \hat{B}_{\mu x} &= B_\mu, \\
\hat{g}_{x\mu} &= -k^2 A_\mu, & \hat{B}_{\mu\nu} &= B_{\mu\nu} - A_{[\mu} B_{\nu]}, \\
\hat{g}_{\mu\nu} &= g_{\mu\nu} - k^2 A_\mu A_\nu, & \hat{V}_x^a &= \ell^a, \\
\hat{\phi} &= \phi + \frac{1}{2} \log k, & \hat{V}_\mu^a &= V_\mu^a + \ell^a A_\mu, \\
\hat{M}_{ab}^{-1} &= M_{ab}^{-1},
\end{aligned} \tag{2.6}$$

we find for the five-dimensional action

$$\begin{aligned}
\mathcal{L}_{m\text{II}} &= \sqrt{|g|} \left\{ e^{-2\phi} \left[R - 4(\partial\phi)^2 + \frac{1}{12} H_{\mu\nu\rho}(B) H^{\mu\nu\rho}(B) + \frac{1}{8} \text{Tr}(\partial M^{-1} \partial M) \right. \right. \\
&\quad \left. \left. + \frac{1}{k^2} (\partial k)^2 - \frac{1}{4} k^2 F_{\mu\nu}(A) F^{\mu\nu}(A) - \frac{1}{4k^2} F_{\mu\nu}(B) F^{\mu\nu}(B) \right] \right. \\
&\quad \left. - \frac{1}{4} k F_{\mu\nu}^a(V) M_{ab}^{-1} F^{b\mu\nu}(V) + \frac{1}{2k} D_\mu \ell^a M_{ab}^{-1} D^\mu \ell^b - 2k m^a M_{ab}^{-1} m^b \right\} \tag{2.7} \\
&+ \frac{1}{8} \epsilon^{\mu\nu\rho\lambda\sigma} \eta_{ab} \left\{ F_{\mu\nu}^a(V) F_{\rho\lambda}^b(V) B_\sigma + 2D_\mu \ell^a F_{\nu\rho}^b(V) B_{\lambda\sigma} - 2D_\mu \ell^a F_{\nu\rho}^b(V) A_\lambda B_\sigma \right. \\
&\quad \left. + 4m^a F_{\mu\nu}^b(V) B_{\rho\lambda} B_\sigma + 2m^a D_\mu \ell^b B_{\nu\rho} B_{\lambda\sigma} \right. \\
&\quad \left. - 4m^a D_\mu \ell^b B_{\nu\rho} A_\lambda B_\sigma + 4m^a m^b B_{\mu\nu} B_{\rho\lambda} B_\sigma \right\}.
\end{aligned}$$

Now the field strengths are given by

$$\begin{aligned}
F_{\mu\nu}(A) &= 2\partial_{[\mu} A_{\nu]}, & F_{\mu\nu}(B) &= 2\partial_{[\mu} B_{\nu]}, \\
H_{\mu\nu\rho}(B) &= 3\partial_{[\mu} B_{\nu\rho]} - \frac{3}{2} A_{[\mu} F_{\nu\rho]}(B) - \frac{3}{2} B_{[\mu} F_{\nu\rho]}(A) \\
F_{\mu\nu}^a(V) &= 2\partial_{[\mu} V_{\nu]}^a + 2m^a B_{\mu\nu} + 2m^a A_{[\mu} B_{\nu]} + 2\ell^a \partial_{[\mu} A_{\nu]}, \\
D_\mu \ell^a &= \partial_\mu \ell^a + 2m^a B_\mu,
\end{aligned} \tag{2.8}$$

and can easily be checked to be invariant under the five-dimensional gauge transformations

$$\begin{aligned}
\delta A_\mu &= \partial_\mu \xi, & \delta B_\mu &= \partial_\mu \Sigma, \\
\delta B_{\mu\nu} &= \partial_{[\mu} \Sigma_{\nu]} + A_{[\mu} \partial_{\nu]} \Sigma + B_{[\mu} \partial_{\nu]} \xi, \\
\delta V_\mu^a &= \partial_\mu \chi^a - m^a \Sigma_\mu + 2m^a \Sigma A_\mu \\
\delta \ell^a &= -2m^a \Sigma.
\end{aligned} \tag{2.9}$$

²Here we denote six-dimensional fields as hatted and five-dimensional fields as unhatted.

Note that the symmetry group of the reduced action is $O(4, 20)$, instead of an $O(5, 21)$ symmetry expected from the massless case [18]. This is of course due to the massive gauge transformations (2.9), which break the $O(5, 21)$ into $O(4, 20)$, by making V_μ^a transform as Stückelberg fields on one hand, while on the other hand the winding vector B_μ becomes massive by eating the ℓ^a 's and the Kaluza-Klein vector A_μ is inert under the Stückelberg transformations.

In the next sections we will perform a Scherk-Schwarz reduction on Type IIB theory compactified over $K3$ and show that it leads to the same action as the one we derived above.

3 Type IIB on $K3$: a review

The six-dimensional Type IIB supergravity is obtained via a compactification of the ten-dimensional Type IIB over $K3$ [19, 20]. Its field content consists of a metric $\hat{g}_{\hat{\mu}\hat{\nu}}$, 5 self-dual and 21 anti-self-dual two-forms $\hat{\mathcal{A}}_{\hat{\mu}\hat{\nu}}^i$ transforming under the vector representation of $O(5, 21)$ and 105 scalars, parametrising an $O(5, 21)/(O(5) \times O(21))$ coset. The scalars can be combined into a symmetric $O(5, 21)$ matrix $\hat{\mathcal{M}}_{ij}^{-1}$.

Due to the (anti-)self-duality conditions, it is impossible to write down a covariant action [21]. However it is consistent to write down a non-self-dual action, dropping the self-duality condition. This non-self-dual action will give the same equations of motion, provided that one inserts the duality condition by hand as an extra equation of motion [22].

The bosonic non-self-dual action is then given by

$$\mathcal{L}_{\text{IIB}} = \sqrt{|\hat{g}|} \left\{ \hat{R} + \frac{1}{8} \text{Tr}(\partial \hat{\mathcal{M}}^{-1} \partial \hat{\mathcal{M}}) + \frac{1}{24} \hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}}^i(\hat{\mathcal{A}}) \hat{\mathcal{M}}_{ij}^{-1} \hat{H}^{j\hat{\mu}\hat{\nu}\hat{\rho}}(\hat{\mathcal{A}}) \right\}, \quad (3.1)$$

where the three-form field strength

$$\hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}}^i(\hat{\mathcal{A}}) = 3\partial_{[\hat{\mu}} \hat{\mathcal{A}}_{\hat{\nu}\hat{\rho}}^i] \quad (3.2)$$

satisfies the (anti-)self-duality condition

$$\hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}}^i(\hat{\mathcal{A}}) = \eta^{ij} \hat{\mathcal{M}}_{jk}^{-1} * \hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}}^k(\hat{\mathcal{A}}). \quad (3.3)$$

Here η^{ij} is the invariant $O(5, 21)$ metric

$$\eta_{ij} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & \eta_{ab} \end{pmatrix}, \quad (3.4)$$

with η_{ab} as defined in (2.3).

In order to write the action (3.1) in the string frame, one has to identify which of the 105 scalars corresponds to the dilaton. This can be done by choosing a particular parametrisation of $\hat{\mathcal{M}}_{ij}^{-1}$, writing the 105 $O(5, 21)$ scalars as 80 scalars contained in an $O(4, 20)$ matrix $\hat{M}_{ab}^{-1}$, 24 scalars $\hat{\ell}^a$ transforming as a vector under $O(4, 20)$ and the dilaton $\hat{\phi}$:

$$\hat{\mathcal{M}}_{ij}^{-1} = \begin{pmatrix} e^{-2\hat{\phi}} + \hat{\ell}^a \hat{M}_{ab}^{-1} \hat{\ell}^b + \frac{1}{4} e^{2\hat{\phi}} \hat{\ell}^4 & -\frac{1}{2} e^{2\hat{\phi}} \hat{\ell}^2 & \hat{\ell}^a \hat{M}_{ab}^{-1} + \frac{1}{2} e^{2\hat{\phi}} \hat{\ell}^2 \hat{\ell}^a \eta_{ab} \\ -\frac{1}{2} e^{2\hat{\phi}} \hat{\ell}^2 & e^{2\hat{\phi}} & -e^{2\hat{\phi}} \hat{\ell}^a \eta_{ab} \\ \hat{M}_{ab}^{-1} \hat{\ell}^b + \frac{1}{2} e^{2\hat{\phi}} \hat{\ell}^2 \eta_{ab} \hat{\ell}^b & -e^{2\hat{\phi}} \eta_{ab} \hat{\ell}^b & \hat{M}_{ab}^{-1} + e^{2\hat{\phi}} \hat{\ell}^c \hat{\ell}^d \eta_{ac} \eta_{bd} \end{pmatrix}, \quad (3.5)$$

where with $\hat{\ell}^2$ we mean $\hat{\ell}^2 = \hat{\ell}^a \hat{\ell}^b \eta_{ab}$.

Hence the action (3.1) in the string frame is give by [23]

$$\mathcal{L}_{\text{IIB}} = \sqrt{|\hat{g}|} \left\{ e^{-2\hat{\phi}} \left[\hat{R} - 4(\partial\hat{\phi})^2 + \frac{1}{8} \text{Tr}(\partial\hat{M}^{-1}\partial\hat{M}) \right] + \frac{1}{2} \partial\hat{\ell}^a \hat{M}_{ab}^{-1} \partial\hat{\ell}^b + \frac{1}{24} \hat{H}^i \hat{\mathcal{M}}_{ij}^{-1} \hat{H}^j \right\}. \quad (3.6)$$

Note that by choosing this particular parametrisation of $\hat{\mathcal{M}}_{ij}^{-1}$, we are forced to write the action (3.6) in a manifest $O(4, 20)$ invariant way, though of course it is still implicit $O(5, 21)$ symmetric. Let us for later convenience also denote the $O(5, 21)$ vector $\hat{\mathcal{A}}_{\hat{\mu}\hat{\nu}}^i$ and its gauge transformations in an explicit $O(4, 20)$ symmetric way:

$$\hat{\mathcal{A}}_{\hat{\mu}\hat{\nu}}^i = \begin{pmatrix} \hat{A}_{\hat{\mu}\hat{\nu}} \\ \hat{C}_{\hat{\mu}\hat{\nu}} \\ \hat{V}_{\hat{\mu}\hat{\nu}}^a \end{pmatrix}, \quad \delta\hat{\mathcal{A}}_{\hat{\mu}\hat{\nu}}^i = \begin{pmatrix} \partial_{[\hat{\mu}} \hat{\xi}_{\hat{\nu}]} \\ \partial_{[\hat{\mu}} \hat{\omega}_{\hat{\nu}]} \\ \partial_{[\hat{\mu}} \hat{\chi}_{\hat{\nu}]}^a \end{pmatrix}. \quad (3.7)$$

4 Scherk-Schwarz reduction of Type IIB

There does not exist a massive version of ten-dimensional Type IIB supergravity, nor is it possible to make a Scherk-Schwarz-like reduction over $K3$. Yet, it is still possible to relate this theory to $m\text{IIB}$ reduced over $K3$, generalising the well known T-duality between massless Type IIA and Type IIB in six dimensions [1, 24]. This can be done by performing a Scherk-Schwarz reduction of six-dimensional Type IIB and mapping it onto the five-dimensional action (2.7) of $m\text{IIB}$. In this way we can construct the massive T-duality rules, completely analogous to the massive T-duality rules between Romans' theory and Type IIB supergravity in ten dimensions [4].

In practice, we will use the techniques explained in [4, 7, 9, 10]. The idea consists in interpreting the five-dimensional Stückelberg transformations (2.9) on the Type IIB side as general coordinate transformations in the compactified x -direction [4]. Giving an extra x -dependence to the fields will generate mass terms and Stückelberg transformations after reduction.

Alternatively, one can also take an $O(5, 21)$ symmetry that shifts the 24 scalars $\hat{\ell}^a \rightarrow \hat{\ell}^a + \Lambda^a$ and gauge this transformation, giving Λ^a a linear x -dependence [10]. The slope parameters in this linear dependence appear as 24 masses after reduction. This particular $O(5, 21)$ transformation that generates the shifts, is given by [25, 10]:

$$\hat{\omega}(\Lambda) = \begin{pmatrix} 1 & 0 & 0 \\ -\frac{1}{2}\Lambda^2 & 1 & \Lambda^a \eta_{ab} \\ -\Lambda^a & 0 & \delta^a_b \end{pmatrix}, \quad (4.1)$$

and acts on the scalars $\hat{\ell}^a$ and the two-forms $\hat{\mathcal{A}}_{\hat{\mu}\hat{\nu}}$ as follows:

$$\begin{aligned} \hat{\ell}^a &\rightarrow \hat{\ell}^a + \Lambda^a \\ \hat{A}_{\hat{\mu}\hat{\nu}} &\rightarrow \hat{A}_{\hat{\mu}\hat{\nu}} \\ \hat{C}_{\hat{\mu}\hat{\nu}} &\rightarrow \hat{C}_{\hat{\mu}\hat{\nu}} + \Lambda^a \hat{V}_{\hat{\mu}\hat{\nu}}^b \eta_{ab} - \frac{1}{2} \Lambda^2 \hat{A}_{\hat{\mu}\hat{\nu}} \\ \hat{V}_{\hat{\mu}\hat{\nu}}^a &\rightarrow \hat{V}_{\hat{\mu}\hat{\nu}}^a - \Lambda^a \hat{A}_{\hat{\mu}\hat{\nu}} \end{aligned} \quad (4.2)$$

In order to match the action (3.6) onto (2.7), we have to chose $\Lambda^a(x) = -2m^a x$ and use the following reduction rules:

$$\begin{aligned}
\hat{g}_{xx} &= -k^{-2} , & \hat{A}_{\mu x} &= A_\mu , \\
\hat{g}_{x\mu} &= -k^{-2} B_\mu , & \hat{A}_{\mu\nu} &= B_{\mu\nu} - B_{[\mu} A_{\nu]} , \\
\hat{g}_{\mu\nu} &= g_{\mu\nu} - k^{-2} B_\mu B_\nu , & \hat{V}_{\mu x}^a &= V_\mu^a + 2m^a x A_\mu , \\
\hat{\phi} &= \phi - \frac{1}{2} \log k , & \hat{V}_{\mu\nu}^a &= V_{\mu\nu}^a - B_{[\mu} V_{\nu]}^a + 2m^a x [B_{\mu\nu} - B_{[\mu} A_{\nu]}] \\
\hat{M}_{ab}^{-1} &= M_{ab}^{-1} , & \hat{C}_{\mu x} &= C_\mu - 2m^a x V_\mu^a \eta_{ab} - 2m^2 x^2 A_\mu \\
\hat{\ell}^a &= \ell^a - 2m^a x & \hat{C}_{\mu\nu} &= C_{\mu\nu} - B_{[\mu} C_{\nu]} - 2m^a x [V_{\mu\nu}^b - B_{[\mu} V_{\nu]}^b] \eta_{ab} \\
& & & - 2m^2 x^2 [B_{\mu\nu} - B_{[\mu} A_{\nu]}]
\end{aligned} \tag{4.3}$$

Note that, apart from the explicit x -dependence, the reduction rules given above vary from the ones in (2.6). In fact we are using the so-called T-dual version of (2.6), in order to map the two actions directly onto the same action in five-dimensions. This technique was used for example in [26].

It is not difficult to see that with this reduction Ansatz the kinetic term for the scalars $\hat{\ell}^a$ reduces in the right way and gives in five dimensions an x -independent result: ³

$$\frac{1}{2} \partial_{\hat{A}} \hat{\ell}^a \hat{M}_{ab}^{-1} \partial^{\hat{A}} \hat{\ell}^b = \frac{1}{2} D_A \ell^a M_{ab}^{-1} D^A \ell^b - 2k^2 m^a M_{ab}^{-1} m^b . \tag{4.4}$$

Note that we generated naturally a covariant derivative for ℓ^a , invariant under the Stückelberg transformations (2.9):

$$D_\mu \ell^a = \partial_\mu \ell^a + 2m^a B_\mu . \tag{4.5}$$

The naive reduction of the field strengths $\hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}}^i(\hat{A})$, however, does give x -dependent results. For example,

$$\hat{H}_{ABX}^a(\hat{V}) = \hat{e}_A^\mu \hat{e}_B^\nu \hat{e}_X^x \left[2\partial_{[\mu} V_{\nu]}^a + 4m^a x \partial_{[\mu} A_{\nu]} + 2m^a B_{\mu\nu} - 2m^a B_{[\mu} A_{\nu]} \right] . \tag{4.6}$$

Hence, instead of identifying $\hat{H}_{ABX}^a(\hat{V})$ with the five-dimensional field strength $F_{AB}^a(V)$, as one does in the case of standard Kaluza-Klein reduction, we define the five-dimensional field strength as

$$\begin{aligned}
k F_{AB}^a(V) &\equiv \hat{H}_{ABX}^a(\hat{V}) + \hat{\ell}^a \hat{H}_{ABX}(\hat{A}) \\
&= e_A^\mu e_B^\nu \left[2\partial_{[\mu} V_{\nu]}^a + 2m^a B_{\mu\nu} - 2m^a B_{[\mu} A_{\nu]} + 2\ell^a \partial_{[\mu} A_{\nu]} \right] .
\end{aligned} \tag{4.7}$$

This definition of $F_{\mu\nu}^a(V)$ is not only x -independent, but also coincides with the definition (2.8) and is therefore invariant under the gauge transformations (2.9). Similarly for the other

³To work out the reduction of the field strengths it is convenient to go to Vielbein notation $\partial_{\hat{A}} \hat{\ell}^a = \hat{e}_{\hat{A}}^{\hat{\mu}} \partial_{\hat{\mu}} \hat{\ell}^a$ and use the following reduction Ansatz: $\hat{e}_{\hat{A}}^{\hat{\mu}} = \begin{pmatrix} e_A^\mu & -B_A \\ 0 & k \end{pmatrix}$.

field strengths:

$$\begin{aligned}
H_{ABC}^a(V) &\equiv \hat{H}_{ABC}^a(\hat{V}) + \hat{\ell}^a \hat{H}_{ABC}(\hat{A}) , \\
kF_{AB}(C) &\equiv \hat{H}_{ABX}^a(\hat{C}) - \hat{\ell}^a \hat{H}_{ABX}^b(\hat{V})\eta_{ab} + \frac{1}{2}\hat{\ell}^a \hat{H}_{ABX}(\hat{A}) , \\
H_{ABC}(C) &\equiv \hat{H}_{ABC}^a(\hat{C}) - \hat{\ell}^a \hat{H}_{ABC}^b(\hat{V})\eta_{ab} + \frac{1}{2}\hat{\ell}^2 \hat{H}_{ABC}(\hat{A}) .
\end{aligned} \tag{4.8}$$

Hence, we find for the five-dimensional field strengths of the reduced Type IIB theory:

$$\begin{aligned}
F_{\mu\nu}(A) &= 2\partial_{[\mu}A_{\nu]} , \\
F_{\mu\nu}(B) &= 2\partial_{[\mu}B_{\nu]} , \\
H_{\mu\nu\rho}(B) &= 3\partial_{[\mu}B_{\nu\rho]} - 3A_{[\mu}\partial_{\nu}B_{\rho]} - 3B_{[\mu}\partial_{\nu}A_{\rho]} , \\
F_{\mu\nu}^a(V) &= 2\partial_{[\mu}V_{\nu]}^a + 2m^a B_{\mu\nu} + 2m^a A_{[\mu}B_{\nu]} + \ell^a F_{\mu\nu}(A) , \\
H_{\mu\nu\rho}^a(V) &= 3\partial_{[\mu}V_{\nu\rho]}^a - 3B_{[\mu}\partial_{\nu}V_{\rho]}^a - 3V_{[\mu}^a\partial_{\nu}B_{\rho]} - 6m^a B_{[\mu}B_{\nu\rho]} + \ell^a H_{\mu\nu\rho}(B) , \\
F_{\mu\nu}(C) &= 2\partial_{[\mu}C_{\nu]} - 2m^a V_{\mu\nu}^b \eta_{ab} + 2m^a B_{[\mu}V_{\nu]}^b \eta_{ab} - \ell^a F_{\mu\nu}^b(V)\eta_{ab} + \frac{1}{2}\ell^2 F_{\mu\nu}(A) , \\
H_{\mu\nu\rho}(C) &= 3\partial_{[\mu}C_{\nu\rho]} - 3C_{[\mu}\partial_{\nu}B_{\rho]} - 3B_{[\mu}\partial_{\nu}C_{\rho]} + 6m^a B_{[\mu}V_{\nu\rho]}^b \eta_{ab} \\
&\quad - \ell^a H_{\mu\nu\rho}^b(V)\eta_{ab} + \frac{1}{2}\ell^2 H_{\mu\nu\rho}(B) .
\end{aligned} \tag{4.9}$$

It can be shown that these field strengths are invariant under the five-dimensional gauge transformations:

$$\begin{aligned}
\delta A_\mu &= \partial_\mu \xi , \quad \delta B_\mu = \partial_\mu \Sigma , \quad \delta \ell^a = -2m^a \Sigma , \\
\delta B_{\mu\nu} &= \partial_{[\mu} \Sigma_{\nu]} + A_{[\mu} \partial_{\nu]} \Sigma + B_{[\mu} \partial_{\nu]} \xi , \\
\delta V_\mu^a &= \partial_\mu \chi^a - m^a \Sigma_\mu + 2m^a \Sigma A_\mu , \\
\delta C_\mu &= \partial_\mu \omega + m^a \chi_\mu^b \eta_{ab} - 2m^a \Sigma V_\mu^b \eta_{ab} , \\
\delta V_{\mu\nu}^a &= \partial_{[\mu} \chi_{\nu]}^a + V_{[\mu}^a \partial_{\nu]} \Sigma + B_{[\mu} \partial_{\nu]} \chi^a + 2m^a \Sigma B_{\mu\nu} - m^a B_{[\mu} \Sigma_{\nu]} , \\
\delta C_{\mu\nu} &= \partial_{[\mu} \omega_{\nu]} + C_{[\mu} \partial_{\nu]} \Sigma + B_{[\mu} \partial_{\nu]} \omega - 2m^a \Sigma V_{\mu\nu}^b \eta_{ab} + m^a B_{[\mu} \chi_{\nu]}^b \eta_{ab} ,
\end{aligned} \tag{4.10}$$

where we have used the following reduction rules for parameters of the the gauge transformations (3.7):

$$\begin{aligned}
\hat{\xi}_x &= 2\xi , \quad \hat{\chi}_x^a = \chi^a + 4m^a x\xi , \quad \hat{\omega}_x = 2\omega - 4m^a x\chi^b \eta_{ab} - 4m^2 x^2 \xi , \\
\hat{\xi}_\mu &= \Sigma_\mu , \quad \hat{\chi}_\mu^a = 2\chi_\mu^a + 2m^a x\Sigma_\mu , \quad \hat{\omega}_\mu = \omega_\mu - 2m^a x\chi_\mu^b \eta_{ab} - 2m^2 x^2 \Sigma_\mu .
\end{aligned} \tag{4.11}$$

With the definitions (4.7)-(4.9) of the five-dimensional field strengths and the parametrisation (3.5) of $\hat{\mathcal{M}}_{ij}^{-1}$, we see that reduction of the kinetic term of the two-forms yields:

$$\begin{aligned}
\hat{H}^i(\hat{\mathcal{A}}) \hat{\mathcal{M}}_{ij}^{-1} \hat{H}^j(\hat{\mathcal{A}}) &= \\
&= e^{-2\phi} \hat{H}^2(\hat{A}) + \hat{M}_{ab}^{-1} \left[\hat{H}^a(\hat{V}) + \hat{\ell}^a \hat{H}(\hat{A}) \right] \left[\hat{H}^b(\hat{V}) + \hat{\ell}^b \hat{H}(\hat{A}) \right] \\
&\quad + e^{2\phi} \left[\hat{H}(\hat{C}) - \hat{\ell}^a \hat{H}^b(\hat{V}) \eta_{ab} + \frac{1}{2} \hat{\ell}^2 \hat{H}(\hat{A}) \right]^2 \\
&= e^{-2\phi} k \left[H^2(B) - 3k^2 F^2(A) \right] + M_{ab}^{-1} \left[H^a(V) H^b(V) - 3k^2 F^a(V) F^b(V) \right] \\
&\quad + e^{2\phi} k^{-1} \left[H^2(C) - 3k^2 F^2(C) \right],
\end{aligned} \tag{4.12}$$

such that the full five-dimensional action (3.6) yields

$$\begin{aligned}
\mathcal{L}'_{m\text{II}} &= \sqrt{|g|} \left\{ e^{-2\phi} \left[R - 4(\partial\phi)^2 + \frac{1}{24} H_{\mu\nu\rho}(B) H^{\mu\nu\rho}(B) + \frac{1}{8} \text{Tr}(\partial M^{-1} \partial M) \right. \right. \\
&\quad \left. \left. + \frac{1}{k^2} (\partial k)^2 - \frac{1}{8} k^2 F_{\mu\nu}(A) F^{\mu\nu}(A) - \frac{1}{4k^2} F_{\mu\nu}(B) F^{\mu\nu}(B) \right] \right. \\
&\quad \left. + \frac{1}{24k} H_{\mu\nu\rho}^a(V) M_{ab}^{-1} H^{b\mu\nu\rho}(V) - \frac{1}{8} k F_{\mu\nu}^a(V) M_{ab}^{-1} F^{b\mu\nu}(V) \right. \\
&\quad \left. + \frac{1}{24k^2} e^{2\phi} H_{\mu\nu\rho}(C) H^{\mu\nu\rho}(C) - \frac{1}{8} e^{2\phi} F_{\mu\nu}(C) F^{\mu\nu}(C) \right. \\
&\quad \left. + \frac{1}{2k} D_\mu \ell^a M_{ab}^{-1} D^\mu \ell^b - 2k m^a M_{ab}^{-1} m^b \right\}.
\end{aligned} \tag{4.13}$$

Note that the Scherk-Schwarz reduction has broken the $O(5, 21)$ symmetry into an $O(4, 20)$, which is of course the group left invariant under the gauging (4.1). Although the group structure of action (4.13) coincides with the one from action (2.7), the actions themselves are clearly different. Yet it turns out that they are equivalent and will become identical upon dualising some of the gauge fields.

As mentioned above, the six-dimensional gauge fields obey the (anti-)self-duality relation (3.3), which forms an essential part of the theory, though not of the action. These self-duality relations have to be taken into account in the reduction to five dimensions, where they give a Hodge duality relation between the two-forms and three-forms. Using the reduction Ansatz (4.7)-(4.9) and the parametrisation (3.5), we find the following five-dimensional duality relations:

$$\begin{aligned}
H^{\mu\nu\rho}(B) &= \frac{1}{2\sqrt{|g|}} e^{2\phi} \epsilon^{\mu\nu\rho\lambda\sigma} F_{\lambda\sigma}(C), \\
H^{\mu\nu\rho}(C) &= -\frac{1}{2\sqrt{|g|}} e^{-2\phi} k^2 \epsilon^{\mu\nu\rho\lambda\sigma} F_{\lambda\sigma}(A), \\
H^{\mu\nu\rho a}(V) &= -\frac{1}{2\sqrt{|g|}} k \eta^{ab} M_{bc}^{-1} \epsilon^{\mu\nu\rho\lambda\sigma} F_{\lambda\sigma}^c(V),
\end{aligned} \tag{4.14}$$

Again we see that the $O(5, 21)$ symmetry of (3.3) gets broken into an $O(4, 20)$ symmetry. Using these duality relations, we can substitute $H_{\mu\nu\rho}^a(V)$, $F_{\mu\nu}(C)$ and $H_{\mu\nu\rho}(C)$ in the action (4.13), leading to

$$\begin{aligned}
\mathcal{L}_{m\text{II}} = & \sqrt{|g|} \left\{ e^{-2\phi} \left[R - 4(\partial\phi)^2 + \frac{1}{12} H_{\mu\nu\rho}(B) H^{\mu\nu\rho}(B) + \frac{1}{8} \text{Tr}(\partial M^{-1} \partial M) \right. \right. \\
& \left. \left. + \frac{1}{k^2} (\partial k)^2 - \frac{1}{4} k^2 F_{\mu\nu}(A) F^{\mu\nu}(A) - \frac{1}{4k^2} F_{\mu\nu}(B) F^{\mu\nu}(B) \right] \right. \\
& \left. - \frac{1}{4} k F_{\mu\nu}^a(V) M_{ab}^{-1} F^{b\mu\nu}(V) + \frac{1}{2k} D_\mu \ell^a M_{ab}^{-1} D^\mu \ell^b - 2k m^a M_{ab}^{-1} m^b \right\} \\
& - \frac{1}{8} \epsilon^{\mu\nu\rho\lambda\sigma} \eta_{ab} \left\{ D_\mu \ell^a F_{\nu\rho}^b(V) (B_{\lambda\sigma} - A_\lambda B_\sigma) + 2(V_\mu^a + \ell^a A_\mu) F_{\nu\rho}^b(V) F_{\lambda\sigma}(B) \right. \\
& \left. - 2(V_\mu^a + \ell^a A_\mu) D_\nu \ell^a H_{\rho\lambda\sigma} \right\}. \quad (4.15)
\end{aligned}$$

After rewriting the topological terms, this action coincides with action (2.7), obtained from dimensionally reducing the massive Type IIA theory.

5 Massive T-duality rules in six dimensions

Now that we reduced two theories $m\text{IIA}$ and IIB in six dimensions onto the same five-dimensional action $m\text{II}$, it is straight forward to construct a direct mapping between the unreduced theories. This mapping is the massive T-duality, a generalisation of the (massless) T-duality between (massless) Type IIA and Type IIB [1, 24, 23].

Comparing the $m\text{IIA}$ and IIB reduction rules (2.6) and (4.3), we find the following $m\text{T}$ -duality rules (denoting the $m\text{IIA}$ fields with a prime):

$$\begin{aligned}
\hat{g}'_{xx} &= 1/\hat{g}_{xx}, & \hat{B}'_{\mu x} &= \hat{g}_{\mu x}/\hat{g}_{xx}, \\
\hat{g}'_{\mu x} &= \hat{A}_{\mu x}/\hat{g}_{xx}, & \hat{B}'_{\mu\nu} &= \hat{A}_{\mu\nu} + 2(\hat{g}_{x[\mu} \hat{A}_{\nu]x})/\hat{g}_{xx}, \\
\hat{g}'_{\mu\nu} &= \hat{g}_{\mu\nu} - (\hat{g}_{\mu x} \hat{g}_{\nu x} - \hat{A}_{\mu x} \hat{A}_{\nu x})/\hat{g}_{xx}, & \hat{V}'^a_x &= \hat{\ell}^a + 2m^a x, \\
\hat{\phi}' &= \hat{\phi} - \frac{1}{2} \log |\hat{g}_{xx}|, & \hat{V}'^a_\mu &= \hat{V}^a_{\mu x} + \hat{\ell}^a \hat{A}_{\mu x}, \\
\hat{M}'^{-1}_{ab} &= \hat{M}^{-1}_{ab},
\end{aligned} \quad (5.1)$$

and inversely (denoting the IIB fields with a prime):

$$\begin{aligned}
\hat{g}'_{xx} &= 1/\hat{g}_{xx}, & \hat{A}'_{\mu x} &= \hat{g}_{\mu x}/\hat{g}_{xx}, \\
\hat{g}'_{\mu x} &= \hat{B}_{\mu x}/\hat{g}_{xx}, & \hat{A}'_{\mu\nu} &= \hat{B}_{\mu\nu} + 2(\hat{g}_{x[\mu} \hat{B}_{\nu]x})/\hat{g}_{xx}, \\
\hat{g}'_{\mu\nu} &= \hat{g}_{\mu\nu} - (\hat{g}_{\mu x} \hat{g}_{\nu x} - \hat{B}_{\mu x} \hat{B}_{\nu x})/\hat{g}_{xx}, & \hat{\ell}'^a &= \hat{V}^a_x - 2m^a x, \\
\hat{\phi}' &= \hat{\phi} - \frac{1}{2} \log |\hat{g}_{xx}|, & \hat{V}'^a_{\mu x} &= \hat{V}^a_\mu - (\hat{V}^a_x - 2m^a x) \hat{g}_{\mu x}/\hat{g}_{xx}, \\
\hat{M}'^{-1}_{ab} &= \hat{M}^{-1}_{ab},
\end{aligned} \quad (5.2)$$

The massive T-duality rules have extra massive corrections with respect to the massless ones in the transformation of the $O(4, 20)$ gauge fields and scalars. Through the Scherk-Schwarz procedure these corrections have a linear dependence on the coordinate in which the T-duality is been performed. Note that we do not give explicitly the T-duality rules relating the $\hat{V}_{\mu\nu}^a$, $\hat{C}_{\mu x}$ and $\hat{C}_{\mu\nu}$ fields to Type IIA fields. It turns out they are non-locally related to $\hat{V}'^a_\mu$, $\hat{g}'_{\mu x}$ and $\hat{B}'_{\mu x}$ respectively, as can be seen from (4.14).

6 Heterotic theory and massive S-duality?

Another well-known duality between $N = 2$ string theories is the string-string duality between the Heterotic string compactified on T^4 and Type IIA compactified on $K3$ [1, 2, 27, 28, 24]. This is a non-perturbative duality, relating the strong coupling regime of one theory to the weak coupling regime of the other and vice versa.

There seems no straightforward generalisation of this duality to a massive string-string duality in six dimensions between massive IIA on $K3$ and Heterotic on T^4 , one of the most obvious reasons being that the symmetry groups do not match. Heterotic on T^4 has an $O(4, 20)$ symmetry and performing a gauging analogous to (4.1) would give a massive five-dimensional theory with $O(3, 19)$ symmetry.

However, Heterotic theory on T^5 has an $O(5, 21)$ symmetry and performing a Scherk-Schwarz reduction by gauging the shift transformation (4.1), gives an $O(4, 20)$ invariant massive $N = 4$ gauged supergravity in four dimensions [10]. Therefore, naively one might think that a massive S-duality might be possible in five dimensions after a Scherk-Schwarz reduction of Heterotic on T^5 and the standard Kaluza-Klein reduction of m II theory, whose action was given in (2.7) and (4.15).

However a closer study reveals that such a massive S-duality does not work, as pointed out in [15]. There are various way to see this:

1. It turns out that, although the two theories have the same field content, the behaviour of the fields under the massive gauge transformations is different in each theory. Where the m II theory (2.7) the $O(4, 20)$ vectors are Stückelberg fields that get eaten by the Kalb-Ramond form, in the Scherk-Schwarz reduction of the Heterotic theory it is the V_μ^a that become massive through the Stückelberg transformation of scalars coming from one of the $O(5, 21)$ vectors [10].
2. Massless string-string-string triality [24] between Heterotic theory on T^4 , Type IIA on $K3$ and Type IIB on $K3$ tells us that the Heterotic theory on T^5 can be thought of as a (standard) Kaluza-Klein reduction of Type IIB. On the other hand, m II theory (2.7) can be thought of as a Scherk-Schwarz reduction of the same Type IIB. Since Scherk-Schwarz and Kaluza-Klein reduction do not commute, it is clear that the Scherk-Schwarz reduction of Heterotic theory on T^5 will give a different result as the Kaluza-Klein reduction of the m II theory.
3. The (massless) string-string-string triality in six dimensions works because in five dimensions there exists only one massless $N = 2$ theory. Hence, all six-dimensional theories can be related to each other after mapping them onto the same five-dimensional

theory. However this is no longer the case for massive theories. Since Scherk-Schwarz and Kaluza-Klein reduction do not commute, the number of inequivalent massive theories grows quickly as one reduces more and more, giving 15, 35 and 71 different theories in six, five and four dimensions respectively [7]. This in itself is of course not a proof for the impossibility of the massive S-duality, but at least it gives an idea of the large number of possibilities, in contrast to the massless case.

7 Conclusions

We generalised the known T-duality rules between six-dimensional Type IIA and Type IIB compactified over $K3$ to massive T-duality between Romans' theory and Type IIB over $K3$. The obtained m T-duality rules receive extra massive correction terms with linear dependence in the direction in which the m T-duality is done. This result is analogous to the ten-dimensional m T-duality rules given in [4].

It was shown in [15] that the six-dimensional massive Type IIA theory interpolates between ten-dimensional Romans' theory and massless Type IIA theory with non-trivial two- and four-form fluxes wrapped over the various two- and four-cycles. For example, using a particular symmetry of the $O(4, 20)$ group, a D8-brane wrapped over the $K3$ can be dualised into a D4-brane solution of massless Type IIA on $K3$ with a four-form flux along the $K3$.

Following the same line of thought, it is not difficult to see that this duality can be extended to include the Type IIB theory: the $O(4, 20)$ group of the five-dimensional massive Type II theory we constructed above will map solutions of Romans' theory into solutions of Type IIB with extra three- and five-form fluxes wrapped over the various non-contractable cycles of $K3 \times S^1$. In particular, the D8-brane wrapped over $K3 \times S^1$ will be dual to a D3-brane solution of Type IIB on $K3 \times S^1$ with an extra F_5 form field in the compactified space.

So far, the generalisations of string dualities between massive supergravities have all been generalisations of the perturbative T-duality. As argued in [15] and section 6, a massive version of the non-perturbative string-string duality is not possible in six-dimensions. The question arises whether massive S-dualities can be constructed for other cases or whether the mass terms turn out to be incompatible with non-perturbative dualities. If the latter turns out to be the case, this adds another question to the mysterious features surrounding (some of) the massive supergravity theories. In fact, the question of the strong coupling behaviour of (some of) these massive supergravity theories is related to the unknown eleven-dimensional origin of Romans' theory and its place in string theory. Still, as Romans' theory should be considered as the real effective low energy limit of Type IIA string theory, being the only supergravity theory that gives rise to D8-brane solutions, one might expect that the U-duality group would, in some way or another, provide a non-perturbative duality for these supergravities.

Note added

Similar results on the six-dimensional massive T-duality rules will also soon appear in [29].

Acknowledgements

The author is very grateful to Dominic Brecher, Clifford Johnson, Mees de Roo, Eduardo Eyras, Laur Järv, Patrick Meessen and Paul Saffin for the useful discussions.

References

- [1] E. Witten, Nucl.Phys. B443 (1995) 85, hep-th/9503124.
- [2] C.M. Hull, P.K. Townsend, Nucl.Phys. B438 (1995) 109, hep-th/9410167.
- [3] J. Polchinski, E. Witten, Nucl.Phys. B460 (1996) 525, hep-th/9510169.
- [4] E. Bergshoeff, M. de Roo, M.B. Green, G. Papadopoulos, P.K. Townsend, Nucl.Phys. B470 (1996) 113, hep-th/9601150.
- [5] L.J. Romans, Phys.Lett. B169 (1986) 374.
- [6] J. Scherk, J.H. Schwarz, Nucl.Phys. B153 (1979) 61.
- [7] P.M. Cowdall, H. Lü, C.N. Pope, K.S. Stelle, P.K. Townsend, Nucl.Phys. B486 (1997) 49, hep-th/9608173.
- [8] H. Lü, C.N. Pope, Mod.Phys.Lett. A12 (1997) 1087, hep-th/9611079.
- [9] I.V. Lavrinenko, H. Lü, C.N. Pope, Nucl.Phys. B492 (1997) 278, hep-th/9611134.
- [10] E. Bergshoeff, M. de Roo, E. Eyras, Phys.Lett. B413 (1997) 70, hep-th/9707130.
- [11] N. Kaloper, R.R. Khuri, R.C. Myers, Phys.Lett. B428 (1998) 297, hep-th/9803066.
- [12] P. Meessen, T. Ortín, Nucl.Phys. B541 (1999) 195, hep-th/9806120.
- [13] N. Kaloper, R.C. Myers, JHEP 05 (1999) 010, hep-th/9901045.
- [14] M. Cvetič, H. Lü, C.N. Pope, Phys.Rev.Lett. 83 (1999) 5226, hep-th/9906221.
- [15] M. Haack, J. Louis, H. Singh, *Massive Type IIA Theory on K3*, hep-th/0102110.
- [16] M.J. Duff, B.E.W. Nilsson, C.N. Pope, Phys.Lett. B129 (1983) 39.
- [17] M.J. Duff, B.E.W. Nilsson, Phys.Lett. B175 (1986) 417.
- [18] J. Maharana, J.H. Schwarz, Nucl.Phys. B390 (1993) 3, hep-th/9207016.
- [19] P.K. Townsend, Phys.Lett. B139 (1984) 283.
- [20] L.J. Romans, Nucl.Phys. B276 (1986) 71.
- [21] M. Marcus, J.H. Schwarz, Phys.Lett. B115 (1982) 111.
- [22] E. Bergshoeff, H.J. Boonstra, T. Ortín, Phys.Rev. D53 (1996) 7206, hep-th/9508091.

- [23] K. Behrndt, E. Bergshoeff, B. Janssen, Nucl.Phys. B467 (1996) 100, hep-th/9512152.
- [24] M.J. Duff, James T. Liu, J. Rahmfeld, Nucl.Phys. B459 (1996) 125, hep-th/9508094.
- [25] E. Bergshoeff, B. Janssen and T. Ortín, Class.Quant.Grav. 13 (1996) 321, hep-th/9506156.
- [26] E. Bergshoeff, C.M. Hull, T. Ortín, Nucl.Phys. B451 (1995) 547, hep-th/9504081.
- [27] A. Sen, Nucl.Phys. B450 (1995) 103, hep-th/9504027.
- [28] J.A. Harvey, A. Strominger, Nucl.Phys. B449 (1995) 535; Erratum-ibid. B458 (1996) 456, hep-th/9504047.
- [29] K. Behrndt, E. Bergshoeff, D. Roest, P. Sundell, in preparation.